



# A sub-pixel resolution enhancement model for multiple-resolution multispectral images

Nicolas Brodu, Dharmendra Singh, Akanksha Garg

## ► To cite this version:

Nicolas Brodu, Dharmendra Singh, Akanksha Garg. A sub-pixel resolution enhancement model for multiple-resolution multispectral images. European Geophysical Union General Assembly 2016, Apr 2016, Vienne, Austria. hal-01287184

**HAL Id: hal-01287184**

**<https://inria.hal.science/hal-01287184>**

Submitted on 25 Aug 2016


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# SUPER-RESOLUTION OF SENTINEL-2 MULTISPECTRAL IMAGES

N. Brodu, Inria Bordeaux, France


Classification of MODIS data with : A. Garg, D. Singh, IIT Roorkee, India



# SENTINEL-2

ESA Satellite, launched in June 2015

13 spectral bands at different resolutions : 60m/pixel, 20m/pixel, 10m/pixel



Goal : put all bands at 10m/pixel



# AVAILABLE TEST DATA (FALL 2015)

R,G,B bands  
at 10m/pixel

Around  
Venice






# RESULT 20M $\rightarrow$ 10M

Example : Near infrared, Band 8A (20m), comparable to Band 8 (10m)

Large band à 10m, narrow spectral band at 20m (targets vegetation)




Original  
Band 8A  
(20m)

# RESULT 20M $\rightarrow$ 10M

Example : Near infrared, Band 8A (20m), comparable to Bande 8 (10m)

Large band à 10m, narrow spectral band at 20m (targets vegetation)




Bande 8A  
résolue  
à 10m

# RÉSULTAT 20M $\rightarrow$ 10M

Example : Near infrared, Band 8A (20m), comparable to Bande 8 (10m)

Large band à 10m, narrow spectral band at 20m (targets vegetation)




Bande 8  
at  
10m

# RÉSULTAT 60M $\rightarrow$ 10M

Example : Visible light, Bande 1 (60m, violet),

To compare with band 2 (10m, blue) ?






# RÉSULTAT 60M $\rightarrow$ 10M

Example : Visible light, Bande 1 (60m, violet),


To compare with band 2 (10m, blue) ?



Super-  
resolved  
Band 1  
at 10m

# RÉSULTAT 60M $\rightarrow$ 10M

Example : Visible light, Bande 1 (60m, violet),  
To compare with band 2 (10m, blue) ?



Original  
Bande 2  
at 10m  
(blue, not  
violet)



Original 60m band 1





Super-resolved 60m band 1





Original 60m band 9



Super-resolved 60m band 9




# METHOD

## Separating color from geometry

- Pixel = mix of different elements
- Mixing information = independent from the spectral band
- Color information information = specific to each band

## Pixel boundaries are arbitrary

- Some information is also shared between nearby neighbors.




Too much pixel variability  
⇒ no long-distance information  
for inferring sub-pixels  
⇒ Local model, ≠ wavelet for ex.


# FOR $20M \rightarrow 10M$

## The problem

Available  
low-resolution  
pixel




To divide in 4  
pixels of higher  
resolution



Constraint:  $\langle H \rangle = L$   
 $\Rightarrow$  3 free paramters  
/ pixel


## The model

Shared values  
between neighbor  
pixels



Depend on  
the spectral  
band


Weights = proportion  
of these shared values  
comprising the pixel



Shared between  
spectral bands  
Constraint:  $\sum w = 1$



# POUR 20M $\rightarrow$ 10M



## Step 1: Fit $S$ , $W$ using the 4 band at 10m

- $H$  available  $\Rightarrow$  1 parameter  $S$  / pixel at 10m is fixed  $\Rightarrow$  no free parameter for  $S$
- 4  $W$  per pixel at 10m, but 3 free parameters and 4 band  $\Rightarrow$  Least squares OK

## Step 2: Find $S$ for the 20m bands, $W$ being fixed (band-independent)

- 2.1 (learning inter-pixels): Fit  $S_L = \sum p_L \times L$ , using neighbors on averaged 10m bands
- 2.2 (apply inter-pixels): Propagate  $S_L = \sum p_L \times L$  on 20m bands
- 2.3 (apply details): Propagate  $S/S_L$  of 10m bands  $\rightarrow$  initial  $S$  value for 20m bands
- 2.4 (apply weights):  $H = W * S$  for 20m bands (+ renormalization  $\langle H \rangle = L$ )

# FOR 60M $\rightarrow$ 10M

Method cannot be applied as such: 36 sub-pixels at 10m / pixel at 60m!


Solution : Intermediate 60m  $\rightarrow$  20m step

- Quick analysis:
  - 9 sub-pixels at 20m, fixed average  $\Rightarrow$  8 free parameters / pixel
  - 6 bands at 20m + 4 at 10m  $\Rightarrow$  10 constraints ( $\approx 9$  with similar B8/B8A)
  - + 3 free parameters per W, shared /10 bandes  $\approx 0.3$  param/pixel $\Rightarrow$  Global least square fit for S,W on 10 bands OK ( $\approx 8.3/9$  param/pixel)
- + Same method for splitting geometry/color : Fit S, W at 20m, then  $S_L$ , then details.

60m original

20m intermediate

10m final



# MODIS DATA


## 2 NASA satellites, cloned

- Visible/infrared
- 2 bands at 250m/pixel, 5 bands à 500m/pixel
- 2 acquisitions / day (final Sentinel-2 scenario = 1 acquisition every 3-4 days)

## Method adaptation

- 2 bands of “high” resolution are not sufficient to fix the mixing weights  $W$
- Hypothesis : The mixing geometry information ( $W$ ) is unchanged over short times
  - ⇒ Fix  $S$ ,  $W$  over several acquisitions (min=2, more are best for clouds)
- E.g. using cloud-free bottom of atmosphere images processed with 16 day data, with overlapping 8-days windows

# EXAMPLE



Original (infrared 1628-1652 nm)




Super-resolution 500m => 250m




# LAND OCCUPATION

Study zone around Roorkee, 200km north of Delhi

classification with original 500m data



with 250 super-resolved data



Blue : water    Purple : urban    Green : Forest    Yellow : Fields    Red : bare soil